

Images in Structural Heart Disease

Superior Mesenteric Artery Embolism Complicating Transcatheter Aortic Valve Replacement

Takehiro Yamashita, MD, PhD^{a,*} , Naoki Iwakiri, MD^a, Eigo Kurebayashi, MD^a, Hitoshi Kimura, MD^b, Masaki Murata, MD^c

^a Department of Cardiology, Cardiovascular Center, Hokkaido Ohno Memorial Hospital, Sapporo, Hokkaido, Japan

^b Department of Surgery, Hokkaido Ohno Memorial Hospital, Sapporo, Hokkaido, Japan

^c Department of Pathology, Hokkaido Ohno Memorial Hospital, Sapporo, Hokkaido, Japan

ARTICLE INFO

Article history:

Submitted 12 May 2021

Revised 11 August 2021

Accepted 3 November 2021

Available online 8 April 2022

Keywords:

Bowel ischemia

Complication

Intravascular aspiration

Superior mesenteric artery embolism

Transcatheter aortic valve replacement

An 83-year-old male was scheduled for transcatheter aortic valve replacement for severe aortic valve stenosis. Although preprocedural computed tomography demonstrated atherosclerotic plaques protruding into the descending aortic lumen (Figure 1), we concluded a transfemoral approach was feasible. However, while advancing the valve system through the descending aorta, the echocardiographer detected a fragment detached from the protruding plaque on transesophageal echocardiography (not recorded). A 26-mm Sapien 3 (Edwards Lifesciences, Irvine, California) was successfully implanted (Figure 2) while transesophageal echocardiography delineated a contrasting configuration of the plaque (Figure 3a and b, Supplemental Video 1). Emergency superior mesenteric artery angiography demonstrated an embolic partial obstruction (Figure 4a), and an



Figure 1. Preprocedural computed tomography showing protruding plaques into the descending aortic lumen.

intravascular aspiration procedure, using an Export Advance catheter (Medtronic, Minneapolis, Minnesota), restored blood flow to the ileocolic territory (Figure 4b). His post-transcatheter aortic valve replacement course

* Address correspondence to: Takehiro Yamashita, MD, PhD, Cardiology, Cardiovascular Center, Hokkaido Ohno Memorial Hospital, 2-1-16-1, Miyanosawa, Nishiku, Sapporo, Hokkaido 063-0052, Japan

E-mail address: t.yamashita@cvc-ohno.or.jp (T. Yamashita).




Figure 2. Aortography after Sapien 3 implantation.




Figure 3. Transesophageal echocardiography (TEE) showing a guidewire (arrow) with the protruding plaque (arrowhead) before (a) and after (b) device advancement.




Figure 4. Superior mesenteric artery angiography showing embolic partial obstruction of the mid portion before the aspiration procedure (a) and restoration of blood flow to ileocolic territory after the procedure (b). Arrow heads indicate partially improved blood flow to iliocolic territory.




Figure 5. Pathology of the aspirate showing cholesterol clefts (green arrows), foamy macrophages (blue arrows), and calcification (yellow arrow) in low (a) and high (b) magnifications.

was uneventful with no signs of bowel ischemia. Pathology showed the retrieved material was a fragment of atherosclerotic plaque with typical cholesterol clefts and foamy macrophages (Figure 5a and b).

ORCIDiDs

Takehiro Yamashita  <https://orcid.org/0000-0001-5918-4308>

Consent Statement

Patient consent for publication of this report and accompanying images was obtained by the authors.

Funding

The authors have no funding to report.

Disclosure statement

The authors report no conflict of interest.

Supplementary Material

Supplemental data for this article can be accessed on the [publisher's website](#).